Application No.:

10/553,092

Filing Date.:

October 13, 2005

AMENDMENTS TO THE CLAIMS

1. (Currently amended) A biologically active substance-immobilized device, which comprises:

a-base particles each comprising a core particle and an organic compound having two or more hydrophilic groups and, wherein the organic compound is immobilized on the core particle by a chemical bond, and

a biologically active substance bonded to the base particles via the organic compound,

wherein the chemical bond is selected from covalent bond, coordinate bond and ionic bond, and wherein the core particle and the biologically active substance are bonded by a reaction with a functional group selected from the group consisting of carbodiimide group, ester group, carbonate group, epoxy group and oxazoline group.

- 2. (Previously presented) The device according to claim 1, monodispersed in an aqueous medium.
- 3. (Currently amended) The device according to claim 1, wherein the base particles has have an average particle diameter of 0.01 to 100 μm .
- 4. (Currently amended) The device according to claim 1, wherein the base particles has have a spherical or substantially spherical shape.
- 5. (Currently amended) The device according to claim $\frac{14}{2}$, wherein at least one of CV_b ratio and CV_c ratio defined by the following equations is 0.6 to 3.0:

 CV_b ratio = CV_1/CV_3

 CV_c ratio = CV_2/CV_3

 CV_1 = (Standard deviation of core particle diameter/Average core particle diameter) x

 CV_2 = (Standard deviation of base particle diameter/Average base particle diameter) x 100

 CV_3 = (Standard deviation of device diameter/average device particle diameter) x 100.

6. (Cancelled)

100

Application No.: 10/553,092

Filing Date.: October 13, 2005

7. (Previously presented) The device according to claim 1, wherein the organic compound is a compound represented by the following formula:

$$A_{x}-(R-X)_{n}-R-A_{y} \tag{I}$$

wherein A_x and A_y independently represent a segment having a functional group that exhibits hydrophilicity and may be identical or different, R independently represents an organic group of two or more valences, X independently represents carbodiimide group, epoxy group or oxazoline group, and n is an integer of 2 to 80.

- 8. (Previously presented) The device according to claim 1, wherein the biologically active substance is selected from a nucleic acid, protein, hapten and saccharide.
- 9. (Previously presented) The device according to claim 1, which is for detecting or measuring a second biologically active substance contained in a sample by using a specific bond of the biologically active substance and the second biologically active substance in the sample.
- 10. (Previously presented) The device according to claim 1, wherein the biologically active substance is an agent for the apeutic treatment of a disease.
- 11. (Previously presented) The device according to claim 7, wherein n is an integer of 2 to 40.
- 12. (Withdrawn-Currently amended) A method of detecting or measuring a second biologically active substance in a sample comprising the step of binding the second biologically active substance to the biologically active substance bound to the base particles in the device of claim 1.
- 13. (New) The device according to claim 1, wherein the organic compound having two or more hydrophilic groups has water solubility.